

NOTE ON A PAPER BY BERNŠTEĬN,
GEL'FAND AND GEL'FAND ON VERMA MODULES

BY

A. VAN DEN HOMBERGH *)

(Communicated by Prof. T. A. SPRINGER at the meeting of January 26, 1974)

Let \mathfrak{G} denote a complex semisimple Lie algebra and \mathfrak{h} a Cartan subalgebra of \mathfrak{G} with dual \mathfrak{h}^* . In [6] VERMA studied "universal modules with a highest weight" M_λ , where $\lambda \in \mathfrak{h}^*$. (For a precise definition of M_λ see below.) He proved that $M_{s_\alpha(\lambda)} \subset M_\lambda$ if $\lambda - s_\alpha(\lambda) = n\alpha$ for some non-negative integer n , and conjectured that every imbedding $M_\mu \subset M_\lambda$ is a composition of imbeddings of that type. This conjecture was proved by I. N. BERNŠTEĬN, I. M. GEL'FAND and S. I. GEL'FAND in [2], in a rather awkward way; their proof doesn't get any simpler if λ is assumed to be integral. In this note a straightforward fundamentally simple proof of the conjecture is given for integral λ , (this is theorem 3).

Let R denote the root system of \mathfrak{G} with respect to \mathfrak{h} and \mathfrak{G}^α the root space corresponding to a root α . Let S be a fundamental system in R . This induces an ordering in R : $R = R_+ \cup R_-$; we set $\mathfrak{H}_\pm = \sum_{\alpha \in R_\pm} \mathfrak{G}^\alpha$. Let $(,)$ be the inverse Killing form on \mathfrak{h}^* ; then

$$\mathfrak{h}_Z^* = \left\{ \lambda \in \mathfrak{h}^* \mid \frac{2(\lambda, \alpha)}{(\alpha, \alpha)} \in \mathbb{Z}, \text{ for all } \alpha \in R \right\}$$

is the lattice of integral elements in \mathfrak{h}^* . Let $C = \{\lambda \in \mathfrak{h}^* \mid (\lambda, \alpha) > 0 \text{ if } \alpha \in R_+\}$ be the highest Weyl chamber with respect to S and \bar{C} its closure. W denotes the Weyl group of \mathfrak{G} and for $\alpha \in R$, $s_\alpha \in W$ is defined by

$$s_\alpha(\lambda) = \lambda - \frac{2(\lambda, \alpha)}{(\alpha, \alpha)} \alpha \text{ for } \lambda \in \mathfrak{h}^*.$$

For $w \in W$, $l(w)$ denotes the length of w . In [1, Chap. VI, no. 1.6, prop. 17] the following is proved:

(*) if $\alpha \in R_+$ then $l(s_\alpha w) > l(w)$ if and only if $w^{-1}(\alpha) \in R_+$.

For $\alpha \in R_+$, we define $H_\alpha \in \mathfrak{h}$ by

$$\lambda(H_\alpha) = \frac{2(\lambda, \alpha)}{(\alpha, \alpha)} \text{ for all } \lambda \in \mathfrak{h}^*.$$

*) This work was supported by the Netherlands Organisation for the Advancement of Pure Research (Z.W.O.).

It is possible to choose $X_\alpha \in \mathfrak{G}^*$ such that $[X_\alpha, X_{-\alpha}] = H_\alpha$ for all $\alpha \in R$.

Let $U(\mathfrak{G})$ denote the universal enveloping algebra of \mathfrak{G} and $U(\mathfrak{R}_-)$ the subalgebra of $U(\mathfrak{G})$ generated by 1 and \mathfrak{R}_- . Let $\delta = \sum_{\alpha \in R_+} \frac{1}{2}\alpha$; then $\delta \in \mathfrak{h}_{\mathbb{Z}}^*$.

For $\lambda \in \mathfrak{h}^*$ we define I_λ as the left ideal of $U(\mathfrak{G})$ generated by \mathfrak{R}_+ and $\{H - (\lambda - \delta)(H) | H \in \mathfrak{h}\}$. Then $U(\mathfrak{G}) = U(\mathfrak{R}_-) \oplus I_\lambda$. The left $U(\mathfrak{G})$ -module $U(\mathfrak{G})/I_\lambda$ is denoted by M_λ ; M_λ has a highest weight vector $f_\lambda = 1 + I_\lambda$ with weight $\lambda - \delta$.

THEOREM 1 (Verma). Let $\lambda, \mu \in \mathfrak{h}^*$. Only two cases are possible:

- (i) $\text{Hom}_{U(\mathfrak{G})}(M_\mu, M_\lambda) = 0$;
- (ii) $\dim_{\mathbb{C}} \text{Hom}_{U(\mathfrak{G})}(M_\mu, M_\lambda) = 1$; every non-trivial homomorphism is an imbedding. Moreover, there is a $w \in W$ such that $w(\mu) = \lambda$. (For a proof, see [6].)

Let $\mu, \lambda \in \mathfrak{h}^*$ and suppose $u \in U(\mathfrak{R}_-)$, $u \neq 0$ and $I_\mu u \subset I_\lambda$, then the right multiplication with u in $U(\mathfrak{G})$, induces a non-trivial homomorphism $M_\mu \rightarrow M_\lambda$ and hence an imbedding, which is denoted by $r(u)$. If, for some $u \in U(\mathfrak{R}_-)$ uf_λ is \mathfrak{R}_+ -extreme (i.e. annihilated by \mathfrak{R}_+), with weight $\mu - \delta$, then $r(u): M_\mu \rightarrow M_\lambda$ is an imbedding.

LEMMA 1. For all $\alpha \in R$ and k a nonnegative integer, $[X_\alpha, X_{-\alpha}^k] = -kX_{-\alpha}^{k-1}(H_\alpha - k + 1)$. If $\alpha \in S$ and $\lambda \in \mathfrak{h}^*$ are such that $\lambda(H_\alpha)$ is a nonnegative integer, then $X_{-\alpha}^{\lambda(H_\alpha)} f_\lambda$ is \mathfrak{R}_+ -extreme with weight $s_\alpha(\lambda) - \delta$.

PROOF. The first statement follows by induction on k . The second statement immediately follows from the first.

Let $Q_L(\mathfrak{R}_-)$ denote the left division ring of quotients of $U(\mathfrak{R}_-)$; see [3, p. 166] for its definition and existence. By $\mathfrak{P}(Q_L(\mathfrak{R}_-))$ we denote the set of all one-dimensional subspaces of $Q_L(\mathfrak{R}_-)$. We define a multiplication in $\mathfrak{P}(Q_L(\mathfrak{R}_-))$ as follows: $\mathbb{C}X \cdot \mathbb{C}Y = \mathbb{C}X \cdot Y$ for all $X, Y \in Q_L(\mathfrak{R}_-)$, $X \neq 0$, $Y \neq 0$.

THEOREM 2 (Verma). There exists a map $u_L: W \times \mathfrak{h}_{\mathbb{Z}}^* \rightarrow \mathfrak{P}(Q_L(\mathfrak{R}_-))$ with:

- (i) $u_L(s_\alpha, \lambda) = \mathbb{C}X_{-\alpha}^{\lambda(H_\alpha)}$, if $\alpha \in S$;
 - (ii) $u_L(w_1 w_2, \lambda) = u_L(w_1, w_2(\lambda)) \cdot u_L(w_2, \lambda)$.
- If $\lambda \in \mathfrak{h}_{\mathbb{Z}}^*$, and $w \in W$, then $M_{w(\lambda)} \subset M_\lambda$ if and only if $u_L(w, \lambda) \subset U(\mathfrak{R}_-)$. In fact, $\text{Hom}_{U(\mathfrak{G})}(M_{w(\lambda)}, M_\lambda) = r(u_L(w, \lambda))$, if $u_L(w, \lambda) \subset U(\mathfrak{R}_-)$.

PROOF. This theorem is a weak version of theorems 4 and 5 in [6], which are harder to prove but not needed for our purpose.

In the same way it can be proved that there exists a map $u_R: W \times \mathfrak{h}_{\mathbb{Z}}^* \rightarrow \mathfrak{P}(Q_R(\mathfrak{R}_-))$, where $Q_R(\mathfrak{R}_-)$ is the right division ring of quotients, with $u_R(s_\alpha, \lambda) = \mathbb{C}X_{-\alpha}^{\lambda(H_\alpha)}$ for $\alpha \in S$, and $u_R(w_1 w_2, \lambda) = u_R(w_1, w_2(\lambda)) \cdot u_R(w_2, \lambda)$. Then one has also for $w \in W$, $\lambda \in \mathfrak{h}_{\mathbb{Z}}^*$: $M_{w(\lambda)} \subset M_\lambda$ if and only if $u_R(w, \lambda) \subset U(\mathfrak{R}_-)$.

If J is a subset of S , let W_J be the subgroup of W generated by the s_α

with $\alpha \in J$. Let $D_J = \{w \in W \mid l(ws_\alpha) > l(w), \text{ for all } \alpha \in J\}$, then D_J is a set of left coset representatives of W/W_J , see [1, exercise 3 of p. 37].

LEMMA 2 (Steinberg). Let $\lambda \in \mathfrak{h}^*$ and $S(\lambda) = \{\alpha \in S \mid (\lambda, \alpha) = 0\}$. Then $W_{S(\lambda)}$ is the stabilizer of λ in W . Suppose $\lambda \in \bar{C}$, $d \in D_{S(\lambda)}$ and $\alpha \in S$; if $d(\lambda)(H_\alpha) < 0$, then $l(s_\alpha d) < l(d)$ and $s_\alpha d \in D_{S(\lambda)}$.

PROOF. The first statement is proved in [4, lemma 1.18]. If $\lambda \in \bar{C}$, $d \in D_{S(\lambda)}$, $\alpha \in S$ and $d(\lambda)(H_\alpha) < 0$, it follows from (*) that $l(s_\alpha d) < l(d)$ and from the definition of $D_{S(\lambda)}$ that $s_\alpha d \in D_{S(\lambda)}$.

The Bruhat ordering in W is defined as follows: If $w_1, w_2 \in W$, then $w_1 < w_2$ if there exist $\alpha_1, \dots, \alpha_k \in R$, such that $l(w_1 s_{\alpha_1} \dots s_{\alpha_{i-1}}) < l(w_1 s_{\alpha_i} \dots s_{\alpha_k})$ for $i = 1, \dots, k$, and $w_1 s_{\alpha_1} \dots s_{\alpha_k} = w_2$. See [7].

Then $w_1 \leq w_2$ if and only if for every reduced expression $w_2 = s_{\alpha_1} \dots s_{\alpha_l}$, $\alpha_1, \dots, \alpha_l \in S$, there exists a subsequence (i_1, \dots, i_p) of $(1, \dots, l)$ such that $w_1 = s_{\alpha_{i_1}} \dots s_{\alpha_{i_p}}$. See [5, exercise on p. 128] or [7].

Then it is clear that:

(**) if $w_1, w_2 \in W$ with $w_1 < w_2$ and $l(s_\alpha w_1) > l(w_1)$, $l(s_\alpha w_2) > l(w_2)$ for some $\alpha \in S$, one has $s_\alpha w_1 < s_\alpha w_2$.

See also [7] for a more general result.

Let $w_0 \in W$ be the longest element of W . Then $S \ni \alpha \mapsto -w_0(\alpha) \in S$, defines an involutory automorphism of the Dynkin diagram, which corresponds to an involutory automorphism A of \mathfrak{G} , which leaves \mathfrak{K}_- invariant. Then $\theta: \mathfrak{K}_- \rightarrow \mathfrak{K}_-$ defined by $\theta(X) = -A(X)$ for $X \in \mathfrak{K}_-$, is an anti-automorphism. This can be uniquely extended to an anti-automorphism of $U(\mathfrak{K}_-)$, which is again denoted by θ . It then follows from the definition of left and right division ring of quotients, that:

$$Q_L(\mathfrak{K}_-) \ni \frac{a}{b} \mapsto \frac{\theta(a)t}{\theta(b)} \in Q_R(\mathfrak{K}_-)$$

defines an anti-isomorphism of $Q_L(\mathfrak{K}_-)$ and $Q_R(\mathfrak{K}_-)$, which is denoted by θ too.

LEMMA 3. $\theta: Q_L(\mathfrak{K}_-) \rightarrow Q_R(\mathfrak{K}_-)$ maps $u_L(w_0 w w_0, w_0 \lambda)$ into $u_R(w, \lambda)^{-1}$, for all $w \in W$ and $\lambda \in \mathfrak{h}_Z^*$.

PROOF. By induction on $l(w)$.

If $l(w) = 1$, then $w = s_\alpha$, for some $\alpha \in S$. Let $k = \lambda(H_\alpha)$, then:

$$\theta(u_L(w_0 s_\alpha w_0, w_0 \lambda)) = \theta(\mathbf{C} X_{w_0(\alpha)}^{-k}) = \mathbf{C} X_{-\alpha}^{-k} = u_R(s_\alpha, \lambda)^{-1}.$$

If $l(w) > 1$, then there exist $\alpha \in S$ and $w' \in W$, such that $l(w') < l(w)$ and $w' s_\alpha = w$. Suppose the lemma holds for w' and all $\lambda \in \mathfrak{h}_Z^*$; then

$$\begin{aligned} \theta(u_L(w_0 w w_0, w_0 \lambda)) &= \theta(u_L(w_0 w' w_0, w_0 s_\alpha(\lambda))) \cdot u_L(w_0 s_\alpha w_0, w_0 \lambda) = \\ &= \theta(u_L(w_0 s_\alpha w_0, w_0 s_\alpha(\lambda))) \cdot \theta(u_L(w_0 w' w_0, w_0 \lambda)) = u_R(s_\alpha, \lambda)^{-1} \cdot u_R(w', \lambda)^{-1} = \\ &= u_R(w, \lambda)^{-1}. \end{aligned}$$

COROLLARY 1. Let $w \in W$ and $\lambda \in \mathfrak{h}_{\mathbf{Z}}^*$. Then $M_{w(\lambda)} \subset M_\lambda$ if and only if $M_{w_0(\lambda)} \subset M_{w_0 w(\lambda)}$.

PROOF. Clear from lemma 3 and theorem 2; note that $u(w^{-1}, w\lambda) = u(w, \lambda)^{-1}$, for all $w \in W$ and $\lambda \in \mathfrak{h}_{\mathbf{Z}}^*$.

The next lemma is a slight extension of lemma 9 of [2]. (The proof remains the same.)

LEMMA 4. Suppose $M_\mu \subset M_\lambda$ and $\alpha \in S$ such that $\mu(H_\alpha)$ is a non-positive integer, then $M_\mu \subset M_{s_\alpha(\lambda)}$.

PROOF. By lemma 1, there are two possibilities, $M_{s_\alpha(\lambda)} \subset M_\lambda$ or $M_{s_\alpha(\lambda)} \supset M_\lambda$. The latter case is trivial, so we suppose that $M_{s_\alpha(\lambda)} \subset M_\lambda$. Suppose $u \in U(\mathfrak{R}_-)$ is such that uf_λ is \mathfrak{R}_+ -extreme and has weight $\lambda - \delta$. It is possible to write $X_{-\alpha}^k \cdot u$ as $u_1 \cdot X_{-\alpha}^{k_1}$, for some $u_1 \in U(\mathfrak{R}_-)$ and a non-negative integer k_1 , which increases unboundedly with k . Hence $X_{-\alpha}^k \cdot uf_\lambda \in M_{s_\alpha(\lambda)}$, for some k . Suppose k minimal with this property. By lemma 1, we have $X_\alpha \cdot X_{-\alpha}^k uf_\lambda = -k(\mu(H_\alpha) - k) \cdot X_{-\alpha}^{k-1} uf_\lambda$; hence $k = 0$, so $uf_\lambda \in M_{s_\alpha(\lambda)}$.

COROLLARY 2. If $\lambda \in \mathfrak{h}_{\mathbf{Z}}^*$ is such that $\lambda(H_\alpha)$ is a nonnegative integer for some $\alpha \in S$, then $M_{s_\alpha(\mu)} \subset M_\lambda$ if $M_\mu \subset M_\lambda$.

PROOF. If we set $\mu' = w_0(\lambda)$, $\lambda' = w_0(\mu)$ and $\alpha' = -w_0(\alpha)$, then $M_{\mu'} \subset M_{\lambda'}$ by corollary 1. Because $\mu'(H_{\alpha'})$ is a nonpositive integer, it follows then from lemma 4 that $M_{\mu'} \subset M_{s_{\alpha'}(\lambda')}$ and hence from corollary 1 again: $M_{s_{\alpha'}(\mu')} = M_{w_0 s_{\alpha'}(\lambda')} \subset M_{w_0(\mu')} = M_\lambda$.

THEOREM 3. Suppose $\lambda \in \mathfrak{h}_{\mathbf{Z}}^* \cap \bar{C}$ and $d, d' \in D_{S(\lambda)}$. If $M_{d(\lambda)} \subset M_{d'(\lambda)}$, then $d \geq d'$.

PROOF. If $l(d) = 0$, then $d = 1$; hence it follows from the fact that $d'(\lambda) - \lambda$ is a sum of positive roots and [1, Chap. VI, no. 1.6, prop. 18] that $d'(\lambda) = \lambda$, and hence that $d' = 1$; see [1, exercise 3 on p. 37]. Now, let $d \neq 1$ and suppose that $M_{d''(\lambda)} \subset M_{d'(\lambda)}$ implies $d'' \geq d'$ if $d'', d' \in D_{S(\lambda)}$ and $l(d'') < l(d)$. Because $l(d) \neq 0$, there is an $\alpha \in S$ such that $l(s_\alpha d) < l(d)$, so $d'' = s_\alpha d < d$ and $d'' \in D_{S(\lambda)}$. By lemma 4 we then have: $M_{d(\lambda)} \subset M_{d'(\lambda)}$ and $M_{d(\lambda)} \subset M_{s_\alpha d'(\lambda)}$. If $d'(\lambda)(H_\alpha) \geq 0$, it follows by corollary 2 that $M_{d''(\lambda)} \subset M_{d'(\lambda)}$ and by induction: $d'' \geq d'$. Hence $d \geq d'$.

If $d'(\lambda)(H_\alpha) < 0$, then by lemma 2: $s_\alpha d' < d'$ and $s_\alpha d' \in D_{S(\lambda)}$. By corollary 2 and the fact that $s_\alpha d'(\lambda)(H_\alpha) = -d'(\lambda)(H_\alpha) > 0$, it follows that $M_{d''(\lambda)} \subset M_{s_\alpha d'(\lambda)}$; hence it follows by induction that $d'' \geq s_\alpha d'$ and by (**) that $d \geq d'$.

BIBLIOGRAPHY

1. BOURBAKI, N., Groupes et Algèbres de Lie. Hermann (Paris).
2. BERNSTEIN, I. N., I. M. GEL'FAND and S. I. GEL'FAND, Structure of representations generated by vectors of highest weight. Funct. An. and Appl. 5, 1-8 (1971).

3. JACOBSON, N., Lie Algebras. Interscience.
4. STEINBERG, R., Endomorphisms of linear algebraic groups. Mem. Amer. Math. Soc. no. 80.
5. ———, Lecture Notes. Yale University (1968).
6. VERMA, D.-N., Structure of certain induced representations of complex semi-simple Lie algebras. Bull. Amer. Math. Soc. 74, 160–166 (1968).
7. ———, A strengthening of the exchange property of Coxeter groups, preprint (1972).